



Original Article

Content Available online at: <https://bpasjournals.com/math-and-stat/>

A New Dimension of Balanced Universally one Sided Design and its Satisfying Optimal Neighbour Condition

K. Praphullo Singh

Author's Affiliation:

Department of Statistics, Thambal Marik College, Oinam, P.O. Nambol, Manipur-795134, India

***Corresponding Author: K. Praphullo Singh**, Department of Statistics, Thambal Marik College, Oinam, P.O. Nambol, Manipur-795134, India
E-mail: kpraphullo@rediffmail.com

How to cite this article: Singh K.P. (2022). A New Dimension of Balanced Universally one Sided Design and its Satisfying Optimal Neighbour Condition. *Bull. Pure Appl. Sci. Sect. E Math. Stat.* 41E(2), 147-152.

ABSTRACT

In an agricultural design of experiment the treatment received by a plot may affect the other response on the neighbouring plots of a same block or it may happen to affect the response on the following plot. For example of the second condition, the tall varieties may affect the other crops grown on the neighbouring plots by their shades. Bailey (2003) has developed such design concerned with the study of one sided neighbour effect, under the above mentioned second condition. This paper gives a new series of Universally Optimal One-Sided Circular Neighbour Balanced designs.

KEYWORDS: One Sided Circular Neighbour Balanced design, Universally Optimal, Equivalence class of sequences.

AMS subject classification: 62K 05, 62K 10 (Primary).

1. Introduction and preliminaries

In agricultural design of experiment where the treatment applied to one experimental plot may affect the response on the neighbouring plots and response on the plot to which, it is applied, Bailey *op.cit.* had proposed a particular type of designs for one sided neighbour effect which are concerned with the study of one-sided neighbour effect only. Such as the applications in the case of sunflower crops as well as in the case of cereal crops where tall varieties may shade the plot on their neighbouring and affect the response of the plot. Similarly, as in the case of pesticide or fungicide experiment where some portion of the treatment applied may spread to the neighbouring plot immediately down wind and spores from the untreated plots, one-sided neighbour effect of the preceding plot-treatment may occur to the following plot. The linear ridge is the form of the blocks of such design (Welham *et al.* (1996)) the design where the plots are in 1-dimension and 2-dimension are studied. Azais, Bailey and Monod (1993) give a catalogue of circular neighbour balanced designs with $t-1$ blocks of size t or t blocks of size $t-1$, where t is the number of treatments. Many contributions are available in the literatures of Smart *et al.* (1994), Langton (1990), David and Kempton (1996). Later on Bailey and Druilhet (2004) have extended the work of Bailey *op. cit.* taking into account the effect of the treatment on the proceeding plot and following plot that on the concerned plot and the effect of the block. Under the name of Circular Neighbour Balanced design, both of such 1-dimensional and 2-dimensional designs are studied. Kumam and Meitei (2006) have contributed a construction-method of such design. For the reference to the text of the paper, let us associate with some few definitions.

Definition: One Sided Circular Neighbour Balanced design is an arrangement of v treatments in b linear blocks of size k (not necessarily distinct) such that (i) each treatment is replicated r times, (ii) every pair of distinct treatments has concurrence μ and (iii) every treatment is followed by each other treatment λ times assuming that in every block the last plot is followed by the first plot. It is denoted by One Sided Circular Neighbour Balanced design $(v, b, r, k, \mu, \lambda)$.

Such design is neighbour balanced as every treatment is followed by each other treatment λ times and also pairwise balanced in the sense that every pair of distinct treatments has concurrence μ . Clearly, $vr = bk$. These designs become circular, after having recommended having a border plot before the first plot of each block, assuming that the treatment already applied to the last plot is applied to this border plot. But its response is not measured. It is only to get the neighbour effect of the treatment in the border plot to the last plot. So, in practical point of view, for conducting an experiment based on such designs of block size k , the planning of the design compels blocks to be of size $k+1$. It becomes Universally Optimal for estimation of the total effect (Bailey and Druihet (2004) proposition 9, page 1657) if

- (i) there are only s different type of treatments in every block,
- (ii) out of s different type of treatments n_1 repeat m times in the block and each of the remaining n_2 occurs $m+1$ times, where $n_1 + n_2 = s$ and
- (iii) all the occurrences of a treatment in a block must be in a single sequence of adjacent plots (possibly including both the last plot and the first plot).

For the future use in the sequel, Universally Optimal One Sided Circular Neighbour Balanced design $(v, b, r, k, s, m, n_1, n_2, \mu, \lambda)$ denotes the design. Clearly, $n_1 = s - n_2$, $n_2 = k - sm$, $bs = v(v-1)\lambda$. The contribution of each block to the sum of the concurrences of all possible pair of treatments is $\theta/2$ where

$$\begin{aligned}\theta &= n_1(n_1-1)m^2 + n_2(n_2-1)(m+1)^2 + 2n_1n_2m(m+1) \\ &= sm(m+1) + k(k-2m-1).\end{aligned}$$

Thus, $b\theta = v(v-1)\mu$. If a class Δ of competing designs contains a design D such that the information matrix C_D is completely symmetric and $\text{trace}(C_D) \geq \text{trace}(C_d)$ for all $d \in \Delta$, then the design D is said to be Universally Optimal. Two sequences of treatments on a block are equivalent if one sequence can be obtained from the other one by releveling the treatments if we denote ξ the equivalence class of the sequence l on the block u of the design d the trace of C_d is given by (Bailey and Druihet *op. cit.*)

$$C(\xi) = \text{trace}(C_{du}).$$

$$= (k - \frac{2}{k} \sum_{i=1}^v g_i^2 + \sum_{i=1}^v h_i) / 2, \text{ where } g_i \text{ is the number of occurrences of treatment } i \text{ in the sequence } l \text{ and } h_i \text{ is}$$

the number of time treatment i is on the left hand side of itself in sequence l .

2. Construction

In this paper as a new dimension in the sphere of construction of Universally Optimal One Sided Circular Neighbour Balanced design, a lemma of Praphullo and Meitei (2006), using difference sets will be recalled such hereafter. Given a set S of size k i.e. $\{i_1, i_2, \dots, i_k\}$, the forward and the backward differences arising from this set are defined as follows:

$$F = (i_2 - i_1, i_3 - i_2, \dots, i_k - i_{k-1}, i_1 - i_k) \text{ and}$$

$$B = (i_1 - i_2, i_2 - i_3, \dots, i_{k-1} - i_k, i_k - i_1) \text{ respectively. Clearly, } B_k = -F_k.$$

2.1. Lemma: Let M be a module of v elements. Consider t be the initial blocks each containing k elements (not necessarily distinct) of M . These t blocks when developed module v generate an Universally Optimal One Sided Circular Neighbour Balanced design with the parameters $v, b = tv, r = kt, k, s, m, n_1, n_2, \mu, \lambda$, if the following conditions are satisfied:

- (i) there are only s different types of treatments in every initial block,

- (ii) each of n_1 out of these s treatments occurs m times and each of the remaining $s-n_1=n_2$ (say) occurs $(m+1)$ times in the block,
- (iii) all the occurrences of a treatment in every initial block is in a single sequence of adjacent plots (assuming the last plot and the first plot are neighbour),
- (iv) among the totality of forward (or backward) differences arising from the t initial blocks, every non-zero element of M occurs exactly λ times,
- (v) among the totality of differences arising from the t initial blocks, every non-zero element of M occurs exactly μ times.

Consider for v ($v=6t+1$, prime), the Galois field, $GF(v)$ exists since v is prime or prime power.

Let x be the primitive element of $GF(v)$. Then $x^{v-1}=x^0=1$ i.e. $x^{6t}=1$,
i.e. $x^{6t}-1=0$, i.e. $(x^{3t}+1)(x^{3t}-1)=0$, i.e. $x^{3t}+1=0$ i.e. $x^{3t}=-1$ since x is primitive

element. Hence $x^{3t}+1=0$ i.e. $x^{3t}=-1$. Therefore, $(x^t+1)(x^{2t}-x^t+1)=0$ Now,

$x^t+1 \neq 0$, for if $x^t+1=0$ then $x^{2t}=1$, which is not possible as x is a primitive element.

Thus, $(x^{2t}-x^t+1)=0$ i.e. $x^{2t}+1=x^t$.

Again, consider now the two sets C_i and C_i^* initial blocks as given below:

$$C_i = \{x^i, x^{2t+i}, x^{4t+i}\} \quad \dots (2.1)$$

$$C_i^* = \{x^{4t+i}, x^{2t+i}, x^i\}, \quad i=0, 1, 2, \dots, (t-1) \quad \dots (2.2)$$

Let, $x^{2t}-1=x^q$.

The six differences arisen from a typical initial blocks are shown below:

From the set (2.1) i.e. $\pm(x^{2t+i}-x^i)$, $\pm(x^{4t+i}-x^i)$, $\pm(x^{4t+i}-x^{2t+i})$.

i.e. $\pm x^i(x^{2t}-1)$, $\pm x^i(x^{4t}-1)$, $\pm x^i(x^{4t}-x^{2t})$.

i.e. $\pm x^i(x^{2t}-1)$, $\pm x^i(x^{2t}+1)(x^{2t}-1)$, $\pm x^i x^{2t}(x^{2t}-1)$.

i.e. $\pm x^{q+i}$, $\pm x^{t+q+i}$, $\pm x^{2t+q+i}$ (2.3)

All positive terms from the set (2.3) are given as

$$x^{q+i}, x^{t+q+i}, x^{2t+q+i}. \quad \dots (2.4)$$

And all the negative terms from the set (2.3) are given as

$$x^{3t+q+i}, x^{4t+q+i}, x^{5t+q+i} \quad \dots (2.5)$$

The differences can be rearranged from the set (2.4) and the set (2.5) as given below,

$$x^{q+i}, x^{t+q+i}, x^{2t+q+i}, x^{3t+q+i}, x^{4t+q+i}, x^{5t+q+i} \quad \dots (2.6)$$

Similarly, from the set (2.2)

$$x^{q+i}, x^{t+q+i}, x^{2t+q+i}, x^{3t+q+i}, x^{4t+q+i}, x^{5t+q+i} \dots (2.7)$$

Remembering that $x^{6t}=1$. We see that among the differences arising out of the two initial blocks, every non-zero elements of $GF(6t+1)$ is repeated exactly twice. It is learnt that every non-zero elements of $GF(6t+1)$ occur exactly once in the set (2.1), and also once in the set (2.2) as $i=0, 1, 2, \dots, (t-1)$, thus among the totality of all possible differences arisen from C_i and C_i^* every non-zero element of $GF(6t+1)$ exactly occurs twice.

The forward differences arisen from the set (2.1) are given at below :

$$\begin{aligned} &\text{i.e. } (x^{2t+i} - x^i), (x^{4t+i} - x^{2t+i}), (x^i - x^{4t+i}), \\ &\text{i.e. } x^i(x^{2t} - 1), x^i(x^{4t} - x^{2t}), (-1)x^i(x^{4t} - 1), \\ &\text{i.e. } x^i(x^{2t} - 1), x^i \cdot x^{2t} \cdot (x^{2t} - 1), (-1)x^i(x^{2t} + 1)(x^{2t} - 1), \\ &\text{i.e. } x^{q+i}, x^{2t+q+i}, x^{3t}, x^i, x^t, x^q. \\ &\text{i.e. } x^{q+i}, x^{2t+q+i}, x^{4t+q+i}, \dots (2.8) \end{aligned}$$

Similarly, the forward difference arisen from the set (2.2) as given below,

$$\text{i.e. } x^{5t+q+i}, x^{3t+q+i}, x^{t+q+i}. \dots (2.9)$$

Combining the forward differences arisen from the set (2.8) and the set (2.9) are given at below.

$$x^{q+i}, x^{t+q+i}, x^{2t+q+i}, x^{3t+q+i}, x^{4t+q+i}, x^{5t+q+i} \dots (2.10)$$

As, the forward differences are of even (odd) and odd (even) power of x for even (odd) number q respectively, and $i=0, 1, \dots, (t-1)$, all the forward differences are all the $6t$ non-zero elements of the Galois field. Thus among the totality of the forward differences arising out of the two sets C_i and C_i^* , every non-zero elements of $GF(v=6t+1)$ occurs exactly once. Developing the initial sets C_i and C_i^* under the reduction modulo of v i.e. $6t+1$, a Universally Optimal One Sided Neighbour Effect design given in the following theorem can be constructed.

2.1. Theorem:

For v ($v=6t+1$, for some t), the construction of an Universally One Sided Neighbour Effect design with the following parameters, is always guarantee $v=(6t+1)$, $b=2t(6t+1)$, $r=6t$, $k=3=s$, $m=1$, $n_1=3$, $n_2=0$, $\mu=2$, $\lambda=1$,

Proof: As all the $3t$ elements in C_i and C_i^* are distinct, $k=s=3t=n_1$, $n_2=0$, $m=1$. An illustration for a construction of Universally Optimal One Sided Neighbour Effect design is made here below.

2.1. Example:

An illustrative example for a construction of Universally Optimal One Sided Neighbour Effect design with parameters $v=(6t+1)$,

$$b=2t(6t+1), r=6t, k=3=s, m=1, n_1=3, n_2=0, \mu=2, \lambda=1,$$

For, $t=2$, the corresponding parameters is

$v=13, b=52, r=12, k=3=s, m=1, n_1=3, n_2=0, \mu=2, \lambda=1$.

From the initial block, $i=0, 1, \dots, (t-1)$, i can take 0, 1, only,

$\{x^i, x^{2t+i}, x^{4t+i}\}$ when, $i=0$, it can make C_0 and C_0^* , and when $i=1$, then C_1 and C_1^* , can be shown as below by developing C_0, C_0^* and C_1, C_1^* under the reduction modulo of 13, since primitive elements of 13 is 2 when, $i=0$, in the set (2.1); and $i=0$, in the set (2.2)

$C_0 = \{x^0, x^{2t+0}, x^{4t+0}\}; C_0^* = \{x^{4t+0}, x^{2t+0}, x^0\}$ again similarly when, $i=1$, in the set (2.1), $C_1 = \{x^1, x^{2t+1}, x^{4t+1}\}$ and $i=1$ in the set (2.2), $C_1^* = \{x^{4t+1}, x^{2t+1}, x^1\};$

under the reduction modulo of 13, a solution of an Universally Optimal One Sided Neighbour Effect design with parameters, $v=13, b=52, r=12, k=3=s=n_1=3, m=1, n_2=0, \mu=2, \lambda=1$

Since the primitive elements of 13 is 2, then the corresponding value of C_0, C_0^* ,

C_1, C_1^* , may be put in the form

$$C_0 = \{x^0, x^{2t+0}, x^{4t+0}\} = \{2^0, 2^4, 2^8\} = \{1, 3, 9\}$$

$$C_0^* = \{x^{4t+0}, x^{2t+0}, x^0\} = \{2^8, 2^4, 2^0\} = \{9, 3, 1\}$$

$$C_1 = \{x^1, x^{2t+1}, x^{4t+1}\} = \{2^1, 2^5, 2^9\} = \{2, 6, 5\}$$

$$C_1^* = \{x^{4t+1}, x^{2t+1}, x^1\} = \{2^9, 2^5, 2^1\} = \{5, 6, 2\}$$

A construction of design may be put as follows by using the developing method.

$\{1, 3, 9\}, \{9, 3, 1\}, \{2, 6, 5\}, \{5, 6, 2\}$
 $\{2, 4, 10\}, \{10, 4, 2\}, \{3, 7, 6\}, \{6, 7, 3\}$
 $\{3, 5, 11\}, \{11, 5, 3\}, \{4, 8, 7\}, \{7, 8, 4\}$
 $\{4, 6, 12\}, \{12, 6, 4\}, \{5, 9, 8\}, \{8, 9, 5\}$
 $\{5, 7, 0\}, \{0, 7, 5\}, \{6, 10, 9\}, \{9, 10, 6\}$
 $\{6, 8, 1\}, \{1, 8, 6\}, \{7, 11, 10\}, \{10, 11, 7\}$
 $\{7, 9, 2\}, \{2, 9, 7\}, \{8, 12, 11\}, \{11, 12, 8\}$
 $\{8, 10, 3\}, \{3, 10, 8\}, \{9, 0, 12\}, \{12, 0, 9\}$
 $\{9, 11, 4\}, \{4, 11, 9\}, \{10, 1, 0\}, \{0, 1, 10\}$
 $\{10, 12, 5\}, \{5, 12, 10\}, \{11, 2, 1\}, \{1, 2, 11\}$
 $\{11, 0, 6\}, \{6, 0, 11\}, \{12, 3, 2\}, \{2, 3, 12\}$
 $\{12, 1, 7\}, \{7, 1, 12\}, \{0, 4, 3\}, \{3, 4, 0\}$
 $\{0, 2, 8\}, \{8, 2, 0\}, \{1, 5, 4\}, \{4, 5, 1\}$

REFERENCES

- [1] Azais, J. M., Bailey, R.A. and Monod. H. (1993). A catalogue of neighbour designs with border plots efficient, *Biometrics*, 49, 1252-1261.
- [2] Bailey, R. A. (2003). Designs for ONE-sided Neighbour Effect, *Jour. Ind. Soc. Ag. Statistics*, 56(3), 302-314.

- [3] Bailey, R.A. and Druilhet, P. (2004). Optimality of neighbour balanced designs for total effects, *Ann. Statist.*, 32, 1650-1661.
- [4] David, O. and Kempton, R. A. (1996). Designs for interference, *Biometrics*, 52, 597-606.
- [5] Kumam, P. and Meitei, K.K. Singh. (2006). A contribution to Universally Optimal One Sided Neighbour Effect design, a contributed seminar paper to *the International Conference On Statistics and Informatics in Agricultural Research, Indian Society of Agricultural Statistics, N. Delhi*, 27-30, Dec. 60(3), 209.
- [6] Langton, S. (1990). Avoiding edge effects in agro forestry experiments; the use of neighbour- balanced designs and guard areas, *Agro forestry systems*, 12, 173-185.
- [7] Smart, L. E., Blight, M. M. Pickett, J. A. and Pye, B. J. (1994). Development of field strategies in incorporating semiochemicals for the control of pea and bean Weevil (*Sitona Lineatus L.*), *Crop. Protection*, 13, 127-135.
- [8] Welham, S. J.; Baily, R. A; Kinsley, A. E. and Hide, G. A. (1996). Designing experiments to examine competition effects between neighbouring plants in mixed populations, *In: XVIII International Biometric Conference Invited Papers*, 97-105.
