

Change in Entropy of Fermionic Field of BHs of Spin $a^* = +5/2$ & $-5/2$ in XRBs

¹Varsha Kumari, ²Anjana Yadav, ³Anuradha Amar, ⁴Baliram Pd Singh and ⁵Dipo Mahto

Author's Affiliations:	¹ Department of Physics, Indian Institute of Engineering, Science and Technology, Sibpur, Kolkata, India ² Department of Education, A. S. College Deoghar, SKM. University, Dumka, India ³ University Department of Physics, BNMU, Madhepura, India ⁴ Associate professor, Department of Chemistry, BN College, TMBU Bhagalpur, India ⁵ Professor, Department of Physics, BCE, Bhagalpur, India
*Corresponding author:	Dipo Mahto Professor & Head, Dept. of Physics, Bhagalpur College of Engineering, Bhagalpur, India E-mail: dipomahto@hotmail.com
ABSTRACT	The research paper discusses the change in entropy of black holes w. r. t. the change in mass as given by $\delta S / \delta M = 8\pi M(1 - 2\Omega M a^* + a^{*2} / 2 - M\Omega a^{*3})$ for spin parameters $a^* = +5/2$ & $-5/2$ and is used to calculate the possible values of different masses of BHs which concludes that this change in entropy decreases due to co-rotation and increases due to counter rotation.
KEYWORDS	Fermionic fields, Spin Parameters and XRBs

Received on 31.03.2025, Revised on 05.06.2025, Accepted on 12.07.2025

How to cite this article Kumari V., Yadav A., Amar A., Singh BP. and Mahto D. Change in Entropy of Fermionic Field of BHs of Spin $a^* = +5/2$ & $-5/2$ in XRBs. *Bulletin of Pure and Applied Sciences- Physics*, 2025;44D (2): 52-55.

INTRODUCTION

The radiation emitting from black holes is not explained by the classical theory, while the quantum theory explains the emission of radiation from a black body [1]. The basis for the formulation of laws of black holes has been discussed [2]. There are so many discussions have been done regarding the entropy of black holes [3,4,5,6,7,8,9,10,11,12].

The present paper discusses the change in entropy of fermionic fields of BHs w. r.t. the change in mass of spin $a^* = +5/2$ & $-5/2$ and calculates their values in XRBs.

THEORETICAL ASPECTS OF THE WORK

For the present purpose of this work, the concerned model for the change in entropy is used given as [13].

$$\delta S / \delta M = 8\pi M(1 - 2\Omega M a^* + a^{*2} / 2 - M\Omega a^{*3}) \quad M(165M - 33) = 0 \quad (5)$$

The equation (1) is applied for $a^* = +5/2$ and $-5/2$ [14,15], then we have

$$\left(\frac{\delta S}{\delta M}\right)_{+5/2} = -\pi M(165M - 33) \quad (2)$$

$$\left(\frac{\delta S}{\delta M}\right)_{-5/2} = \pi M(165M + 33) \quad (3)$$

The maximum or minimum values of mass can be obtained under the following condition.

$$\left(\frac{\delta S}{\delta M}\right)_{\pm 3/2} = 0 \quad (4)$$

When the equation (1) is applied on the equations (2) and (3), we get..

$$M(165M + 33) = 0 \quad (6)$$

The solutions of the above both equations are given below.

$$M=0 \text{ or } M=33/165 \quad (7)$$

$$M=0 \text{ or } M=-33/165 \quad (8)$$

The solutions for the masses are 0, 33/165 and -33/165. This shows the zero, positive and negative masses respectively. The positive mass explains the gravity. The negative mass provides the naked singularity & dark matter. The zero mass has also existence as explained by the general relativity and quantum theory. Now by the use of proper data, change in entropy w.r.t. the mass for XRBs is listed in the table 1.

Table 1: The change in entropy for $a^* = +5/2$ and $-5/2$.

S.N.	Mass in (M_{\odot})	$\left(\frac{\delta S}{\delta M}\right)_{+5/2} = -\pi M(165M - 33)$ [Joule/Kelvin/kg] $\times 10^{64}$	$\left(\frac{\delta S}{\delta M}\right)_{-5/2} = \pi M(165M + 33)$ [Joule/Kelvin/kg] $\times 10^{64}$
1	5 M_{\odot}	-5 . 1810	5 . 1810
2	6 M_{\odot}	-7 . 4610	7 . 4610
3	7 M_{\odot}	-10 . 1500	10 . 1500
4	8 M_{\odot}	-13 . 2600	13 . 2600
5	9 M_{\odot}	-16 . 7900	16 . 7900
6	10 M_{\odot}	-20 . 7200	20 . 7200
7	11 M_{\odot}	-25 . 0600	25 . 0600
8	12 M_{\odot}	-29 . 8400	29 . 8400
9	13 M_{\odot}	-35 . 0200	35 . 0200
10	14 M_{\odot}	-40 . 6200	40 . 6200
11	15 M_{\odot}	-46 . 6300	46 . 6300
12	16 M_{\odot}	-53 . 0500	53 . 0500
13	17 M_{\odot}	-59 . 8900	59 . 8900
14	18 M_{\odot}	-67 . 1500	67 . 1500
15	19 M_{\odot}	-74 . 8100	74 . 8100
16	20 M_{\odot}	-82 . 900	82 . 900




Figure 1: The variation of entropy change of BHs $a^* = +5/2$ and $-5/2$.

RESULT AND DISCUSSION

The present work starts with consideration of the model for change in entropy given as:

$$\delta S / \delta M = 8\pi M(1 - 2\Omega M a^* + a^{*2} / 2 - M\Omega a^{*3})$$

The observation of above equation indicates that the change in entropy depends on mainly three parameters like angular velocity, spin and mass. The above equation is used for two spins like $a^* = +5/2$ & $-5/2$, two equations(2) & (3) are obtained for discussions and their values listed in the table concludes that change in entropy decreases the entropy for $a^* = +5/2$, while increases the entropy change for $a^* = -5/2$.

For clear conception, the graph is plotted betn the mass of BHs and corresponding change in entropy with change in mass for XRBs with the help of equations (2) & (3). This is shown in the Figs.1. We observe that the variation is symmetrical for either types of spin parameters for co-rotation and counter rotations.

CONCLUSION

The entropy changefor $a^* = +5/2$ showsthe decrease in entropy change, while increases the entropy change for $a^* = -5/2$.The present model

also gives the naked singularity as well as dark matter.

REFERENCES

1. Hawking SW. Particle creation by black holes. **Commun Math Phys.** 1975;43:199–220.
2. Bardeen JM, Carter B, Hawking SW. The four laws of black hole mechanics. **Commun Math Phys.** 1973;31(2):161–170. doi:10.1007/BF01645742.
3. Strominger A, Vafa C. Microscopic origin of Bekenstein–Hawking entropy. **Phys Lett B.** 1996;379(1–4):99–104. doi:10.1016/0370-2693(96)00345-0.
4. Traschen J. An introduction to black hole evaporation. **arXiv preprint.** 2000. arXiv:gr-qc/0010055.
5. Dabholkar A. Black hole entropy in string theory – a window into the quantum structure of gravity. **Curr Sci.** 2005;89(12):2133–2143.
6. Narayan R. Black holes in astrophysics. **New J Phys.** 2005;7:199.
7. Bekenstein JD. Bekenstein–Hawking entropy. **Scholarpedia.** 2008;3(10):7375.

8. Carlip S. Black hole thermodynamics and statistical mechanics. **Lect Notes Phys.** 2009;769:89-123.
9. Mahto D, Kumari K, Sah RK, Singh KM. Study of non-spinning black holes with reference to the change in energy and entropy. **Astrophys Space Sci.** 2012;337:685-691.
10. Mahto D, Singh AK, Ram M, Vineeta K. Change in internal energy and enthalpy of the black holes. **Int J Astrophys Space Sci.** 2013;1(4):61-65.
11. Mahto D, Kumari A. Change in entropy of spinning black holes due to corresponding change in mass in XRBs. **Int J Astron Astrophys.** 2018;8:171-177.
12. Mahto D, Paswn R, Kumari K, Kumar B. Change in entropy of fermionic fields of black holes with respect to mass. **J Inf Comput Sci.** 2020;10(9):342-351.
13. Mahto D, Paswn R, Kumar B, Kumari A. Change in entropy of bosonic fields of black holes with respect to the mass for maximum co-rotation and counter rotation. **J Inf Comput Sci.** 2020;10(9):357-369.
14. Tayal DG. **Nuclear physics.** New Delhi: Himalaya Publishing House; 1990.
15. Yash. An introduction to spin in quantum mechanics (Spin: Explained). **Quantaphy.** 2022
